## Hurricanes and Global Warming: Expectations Versus Observations

KIII 25 September

12 June, 2015

NASA Jet Propulsion Laboratory

Chris Landsea, National Flurricane Center, Miami, USA

Chris.Landsea@noaa.gov



Wilma 7







# Atmospheric & Oceanic Sciences

## Hurricanes and Global Warming: Expectations Versus Observations

KIII 25 September

12 June, 2015

NASA Jet Propulsion Laboratory

Chris Landsea, National Flurricane Center, Miami, USA

Chris.Landsea@noaa.gov



Wilma 7







#### Bloomberg Businessweek IT'SGLOBAL WARMING, STUPID









## Global Warming – Past Temperature Changes

#### Global Land-Ocean Temperature Index



http://data.giss.nasa.gov/gistemp/graphs/Fig.A2.gif


# Nature's great heat engine... The Hurricane




#### Global Warming and Hurricane Winds: Theory and Modeling Work Suggest ~1% Increase Today




### Global Warming and Hurricane Winds: Theory and Modeling Work Suggest ~1% Increase Today




Global Warming and Hurricanes:
Theory and Modeling Work Suggest ~3% wind increase with a DECREASE in frequency by late 21st Century




Global Warming and Hurricanes:
Theory and Modeling Work Suggest ~3% wind increase with a DECREASE in frequency by late 21st Century







#### Tropical Convection Changing with Global Warming




Global Warming and Hurricanes:
Theory and Modeling Work Suggest ~3% wind increase with a DECREASE in frequency by late 21st Century



Global Warming and Hurricanes:
Theory and Modeling Work Suggest ~3% wind increase with a DECREASE in frequency by late 21st Century




#### Increased Wind Shear from Global Warming




Vecchi and Soden (2007)


#### Model projections of atmospheric steering of Sandy-like superstorms

Elizabeth A. Barnes<sup>a,b,1</sup>, Lorenzo M. Polvani<sup>b,c</sup>, and Adam H. Sobel<sup>b,c</sup>





"We demonstrate that climate models consistently project a decrease in the frequency and persistence of the westward flow that led to Sandy's unprecedented track, implying that future atmospheric conditions are less likely than at present to propel storms westward into the coast."






#### U.S. Tropical Storm and Hurricane Damages \$BILLIONS Annually - Inflation Adjusted







#### Hurricane Strikes vs Population for Miami-Dade, Florida



#### NORMALIZED DAMAGE....

Estimated direct damage if past storms made landfall with present-day societal conditions

ND = f(inflation, coastal population, wealth)
Pielke and Landsea (1998)

ND = f(inflation, coastal housing, wealth) Pielke et al. (2008)

### 1926 Great Miami Hurricane

\$165 Billion Today

Miami Beach 1926

Miami Beach 2006







**Wendler Collection** 

Joel Gratz © 2006

U.S. Tropical Storm and Hurricane Damages \$BILLIONS Annually - Normalized





"This record [of Atlantic tropical cyclone counts] .... shows a strong, long-term relationship with tropical **Atlantic August-October** SST...The underlying factor appears to be the influence of (primarily anthropogenic) forced large-scale warming."

Eos, Vol. 87, No. 24, 13 June 2006



#### Atlantic Hurricane Trends Linked to Climate Change


PAGES 233, 238, 241


Increases in key measures of Atlantic hurricane activity over recent decades are believed to reflect, in large part, contemporaneous increases in tropical Atlantic warmth [e.g., Emanuel, 2005]. Some recent studies [e.g., Goldenberg et al., 2001] have attributed these increases to a natural climate cycle


termed the Atlantic Multidecadal Oscillation (AMO), while other studies suggest that climate change may instead be playing the dominant role [Emanuel, 2005; Webster et al., 2005].

Using a formal statistical analysis to separate the estimated influences of anthropogenic climate change from possible natural cyclical influences, this article presents results indicating that anthropogenic factors are likely responsible for long-term trends in tropical Atlantic warmth and tropical cyclone

### Linking Atlantic storm numbers to ocean temperatures








## **RECONNAISSANCE FLIGHT PATH** NOAA **Aircraft** Orion-P3s "ALPHA" Air Force **Pattern** C-130s lispaniola







#### Atlantic Hurricane Database Re-Analysis Project

http://www.aoml.noaa.gov/hrd/data\_sub/re\_anal.html





1851 through 1955 changes accepted and officially adopted by NHC. Revisions for 1956 to 1963 have been submitted and are being considered. Remainder of 20<sup>th</sup> Century currently being reanalyzed.





### RE-ANALYSES NEED TO BE CONDUCTED GLOBALLY!!!













#### **Atlantic Basin Major Hurricanes**


1944 to 2014 (Reanalyzed through 1955)






## Atlantic Multidecadal Oscillation - SSTs, Vertical Shear and Cyclogenesis

Composites around AMM AMM(+) AMM(-)






#### The Recent Increase in Atlantic Hurricane Activity: Causes and Implications

Stanley B. Goldenberg, 1\* Christopher W. Landsea, 1
Alberto M. Mestas-Nuñez, 2 William M. Gray 3


The years 1995 to 2000 experienced the highest level of North Atlantic hurricane activity in the reliable record. Compared with the generally low activity of the previous 24 years (1971 to 1994), the past 6 years have seen a doubling of overall activity for the whole basin, a 2.5-fold increase in major hurricanes (≥50 meters per second), and a fivefold increase in hurricanes affecting the Caribbean. The greater activity results from simultaneous increases in North Atlantic sea-surface temperatures and decreases in vertical wind shear. Because these changes exhibit a multidecadal time scale, the present high level of hurricane activity is likely to persist for an additional ~10 to 40 years. The shift in climate calls for a reevaluation of preparedness and mitigation strategies.






#### **Atlantic Basin Major Hurricanes**


1944 to 2014 (Reanalyzed through 1955)







Warm
Atlantic
Phase
Remains
Thus Far...

Expectation for Active Hurricane Seasons Overall to Continue







## Overall Tropical Storm and Hurricane Changes Due to Global Warming by 2100

Frequency: Numbers may see a moderate decrease (~25%)

Wind Intensity: Small increase (~3% stronger)

**Storm Surge**: Small increase (~3% higher) produced by the hurricane (must also add on additional amount from general sea level rise)

**Rainfall**: Moderate increase per tropical storm and hurricane (~10% within 200 mi of storm), but reduced frequency may offset increases


**Genesis**: Tropical storms and hurricanes to form **slightly farther away** from North America

**Track**: Tropical storms and hurricanes to recurve slightly more often over water and remain away from land

#### **Hurricanes and Global Warming**

Opinion piece by Christopher W. Landsea November 2011





#### **Hurricanes as the Poster-Child for Global Warming**

Hurricanes have been depicted as the literal poster-child of the harmful impacts of global warming. Without argument, hurricanes (which also include storms known as "typhoons" in the Northwest Pacific and "severe tropical cyclones" in the Indian and Southwest Pacific) are extremely destructive and often responsible for the deaths of hundreds and occasionally thousands of people. As an example, Hurricane Katrina was responsible for the death of ~1200 and about \$108 billion in damages. The before and after pictures of the home of David and Kimberly King of Waveland, Mississippi show the incredible power of that hurricane's storm surge and winds.

Opinion Piece on Hurricanes and Global Warming

http://www.aoml.noaa. gov/hrd/Landsea/gw\_ hurricanes/index.html



 Relatively minor manmade hurricane changes decades into the future


- Relatively minor manmade hurricane changes decades into the future
- 2. Huge decadal scale hurricane variations occurring today

- Relatively minor manmade hurricane changes decades into the future
- 2. Huge decadal scale hurricane variations occurring today
- 3. Huge vulnerability issues today, which will get increasingly severe as population grows

- 1. Relatively minor manmade hurricane changes decades into the future
- 2. Huge decadal scale hurricane variations occurring today
- 3. Huge vulnerability issues today, which will get increasingly severe as population grows
- 4. Need to address vulnerability issues now (no need to invoke possible minor changes decades from now):
  - a. Improved hurricane observational network


#### CYGNSS

The Cyclone Global Navigation Satellite System



#### **XOVWM**

The Next-Generation Ocean Surface Vector Winds Mission



Areas of high rain

Fig. 4 Simulated QuikSCAT, DFS and XOVWM wind vector retrievals for Katrina-like hurricane.

- Relatively minor manmade hurricane changes decades into the future
- 2. Huge decadal scale hurricane variations occurring today
- 3. Huge vulnerability issues today, which will get increasingly severe as population grows
- 4. Need to address vulnerability issues now (no need to invoke possible minor changes decades from now):
  - a. Improved hurricane observational network
  - b. Improved hurricane modeling/forecasting
  - c. Improved building codes/land use
  - d. Improved evacuation/shelter plans

# Hurricanes and Global Warming: Expectations Versus Observations

KIII 25 September

12 June, 2015

NASA Jet Propulsion Laboratory

Chris Landsea, National Flurricane Center, Miami, USA


Chris.Landsea@noaa.gov



Wilma 7








Maximum Potential Intensity Change Atlantic Basin: +1% stronger per °C SST change Vecchi and Soden (2007)

## Merging paleotempestology (sediment core records) with modern hurricane data



### Landfalling Strong Hurricanes 1851 to 2009

